

Blue-Green Algae in Wisconsin: Results of a Two-Year Statewide Monitoring Program



Elisabeth Harrahy, Ph.D.

Wisconsin Department of Natural Resources

Dawn A. Karner, Curtis Hedman, William Krick, Steve Geis

Wisconsin State Laboratory of Hygiene

Introduction

- Under certain conditions, blue-green algae can increase in number to “bloom” densities.
- Factors that may be involved include:

- Nutrients

- P; some can fix N
 - recent studies: N just as important

- Temperature

- Water depth

- Wind

- calm, low turbulence
 - gas-filled vesicles (most)

- Zebra mussels

- filter water → deeper pen. sun.
 - eat all except blue-green algae



Monona Golf Course Pond

Introduction

Issues Associated with Blue-Green Algae Blooms

- Discolored water
- Taste and odor problems
- Reduced light penetration
- Dissolved oxygen depletions during die-off
- Toxin production



Introduction

Blue-Green Algal Toxins

- Some species can produce one or more toxins



Red Cedar R.

- Those that can produce toxins do not produce toxins at all times (genetic strain, environmental conditions)
- Toxins can affect
 - Fish and other aquatic life
 - Livestock (reports of deaths date back to 1878)
 - Pets (reports of dog deaths widespread)
 - Humans (70 deaths in Brazil, 1996; 1 in WI, 2002)
- Exposure routes in humans: dermal, oral (water or food), inhalation, dialysis

Dermatotoxins and Gastrointestinal Toxins

- Affect skin and mucous membranes
- Can cause rashes, respiratory illness, headaches, gastrointestinal upset
- lyngbyatoxin, lipopolysaccharide endotoxins

Hepatotoxins

- Affect the liver (cell membrane integrity)
- Can cause hemorrhage, tissue damage, tumors, liver cancer, death
- microcystins, nodularins

Cytotoxins

- Affect the liver and other organs (protein synthesis)
- Can cause chromosome loss, DNA strand breakage, damage to organs
- cylindrospermopsin

Neurotoxins

- Affect the central nervous system (acetylcholinesterase, sodium channels)
- Can cause seizures, paralysis, respiratory failure, death
- anatoxin-a, saxitoxin

Introduction

Guidelines



- Currently no surface water quality criteria, no drinking water standards in U.S. for blue-green algae or for blue-green algal toxins
- World Health Organization guidelines:
 - 100,000 cells/mL = moderate risk to human health
 - 1 $\mu\text{g/L}$ microcystin-LR = drinking water guideline

Background on WI's BGA Monitoring Program

2003

- Limited monitoring conducted for *Cylindrospermopsis* sp.
- Sampled 31 eutrophic lakes
- Cylindro detected in 6



Cylindrospermopsis sp.

2004

- While planning to repeat study received reports of dogs dying (early June)
- High concentrations of *Anabaena* sp. (and anatoxin-a)
- Decided to expand study



Sampling Design



Lakes

5 lakes in each of 5 regions,
5 sample time points



Ponds

8 ponds in SCR only,
5 sample time points

*Most locations selected were **eutrophic** or known to experience blooms;
many were **beaches**; all were **accessible** by the public.
NOT a random sample of lakes and ponds.

Sampling Methods



- Three samples were collected near shore at each location
 - BGA ID and enumeration (plastic, preserved)
 - chlorophyll-a analysis (plastic, in dark, on ice)
 - blue-green algal toxins analysis (amber glass, on ice)
- Shipped overnight to WI State Laboratory of Hygiene


Identification and Enumeration Methods

Tier I Analysis

- all samples
- nanoplankton chamber
- rough estimate in natural units/mL


Tier II Analysis

- select number of samples
- Utermöhl settling chamber
- cells/mL and biovolume (mm^3/m^3)



Analytical Methods

- Blue-green algal toxin analysis was performed by HPLC/MS/MS
- Three toxins in one run
- Report limits were 0.50 µg/L for anatoxin-a and cylindrospermopsin, and 1.00 µg/L for microcystin-LR



Results

2004:

31 lakes	148 samples
10 ponds	38 samples
1 river	1 sample

Total: 187 samples

2005:

38 lakes	154 samples
8 ponds	35 samples
1 river	5 samples

Total: 194 samples

(each site sampled 1 to 6 times)



L. Kegonsa

Results: Blue-Green Algae

2004:

BGA detected in 138/187 samples = 74%

Hot spots: SCR and WCR

2005:

BGA detected in 143/194 samples = 74%

Hot spots: SCR and WCR



L. Menomin
2004
microcystin



Colladay P.
2004 & 2005
anatoxin-a

Results: Toxins



Number of samples analyzed

2004: 45/187 samples = 24%

2005: 34/194 samples = 18%

(Note: selected samples with high concentrations of BGA)

Results: Toxins

2004: 31/45 samples

anatoxin-a	in 5	1.5 – 110 $\mu\text{g/L}$
microcystin-LR	in 26	1.2 – 7,600 $\mu\text{g/L}$

2005: 14/34 samples

anatoxin-a	in 3	0.68 – 2.7 $\mu\text{g/L}$
microcystin-LR	in 13	1.2 – 450 $\mu\text{g/L}$

Guideline Values

anatoxin-a	3 $\mu\text{g/L}$
microcystin-LR	1 $\mu\text{g/L}$



Results: Toxins

Relationship Between Toxins and Blue-Green Algae

- Concentration of microcystin **was not** significantly correlated to the sum concentration of those species of blue-green algae (in natural units/mL, cells/mL, or mm³/m³) capable of producing microcystin (5 species).
- Concentration of microcystin **was** significantly correlated to concentration of *Microcystis* sp.



L. Maria

Results: Chlorophyll-a



-Concentration of Chl-a ranged from 2.26 – 126,000 $\mu\text{g/L}$ (!)


Relationship Between Chl-a and Blue-Green Algae

-In general, concentration of chl-a **was** significantly correlated with concentration of blue-green algae (natural units/mL, cells/mL, or mm^3/m^3), but r^2 values were low.

Results: Chlorophyll-a

Relationship Between Chl-a and Toxins

- Concentration of Chl-a was significantly correlated with concentration of the toxin, microcystin.



2005 $r^2 = 0.31$

Information Sharing




- Program not designed to provide real-time information
- Sent letter and background info to public health officers
- Results shared with local and state public health agencies when Tier I analysis showed sum concentration BGA > 5,000 natural units/mL (33 alerts in 2004; 42 in 2005)
- Only the public health agencies have authority to close or post beaches (responses differed by counties)

Information Sharing

Response by Local Public Health Departments

- Some closed/reopened beaches
- Some posted advisory signs
- One obtained funds to do additional sampling
- All who posted used our generic advisory sign
- Almost all called and asked for more information



Information Sharing

Samples Counted in cells/mL (end of season)

2004:

>5,000 natural units/mL: 22
20 > 100,000 cells/mL

<5,000 natural units/mL: 13
2 > 100,000 cells/mL

2005:

>5,000 natural units/mL: 25
24 > 100,000 cells/mL

<5,000 natural units/mL: 9
1 > 100,000 cells/mL



Research Needs

- Development of threshold values or criteria for blue-green algae and their toxins (WHO values based on *Microcystis* sp. and microcystin-LR toxin concentrations only)
- Development of analytical methods to analyze additional toxins we suspect may be present (e.g., anatoxin-as, BMAA, saxitoxin)
- Studies on the effectiveness of nutrient management plans in reducing the frequency, duration, and severity of blue-green algae blooms
- Studies on the persistence of toxins in water after bloom subsides
- Studies on bioaccumulation of toxins by freshwater fish (BMAA?)

Acknowledgements

Kelsy Anderson
Jesse Baumann
Heidi Bunk
Jim Cahow
Diana Downes
Rachel Gall
Susan Graham
Mark Hazuga
Dan Helsel
Lacey Hoyt
Andy Jegerlehner
Jim Klosiewski
Cindy Koperski



Jim Kreitlow
Paul La Liberte
Michelle Lehner
Trent Miller
Brenner Perryman
Craig Roesler
Ken Schreiber
Lisa Schultz
Greg Searle
Donna Sefton
Mark Sessing
Linda Vogen